

## Heavy Metal Concentrations in Ceiling Fan and Roadside Car park Dust Collected from Residential Colleges in Universiti Putra Malaysia, Serdang, Selangor

C. K. Yap<sup>1\*</sup>, Chew, W. Y.<sup>1</sup>, and S. G. Tan<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Science,

<sup>2</sup>Department of Cell and Molecular Biology,

Faculty of Biotechnology and Biomolecular Science,

Universiti Putra Malaysia, 43400 Serdang,

Selangor, Malaysia

\*E-mail: yapckong@hotmail.com

### ABSTRACT

In this study, dust samples were collected from 4 residential colleges (K2, K5, KOSAS and K10) in Universiti Putra Malaysia (UPM) Serdang, Selangor. The samples were collected from ceiling fans and car parks roadside dust. Sand dust on top of the car park cover were collected using polyethylene brush, tray and kept in a polyethylene bag. Dust from ceiling fan on the first floor was collected and put into a polyethylene bag. The dust samples collected were analysed for the concentrations of Cd, Cu, Fe, Ni and Zn. It was found that K5 [Cu ( $62.94 \pm 0.77$   $\mu\text{g/g dw}$ ), Fe ( $1802.40 \pm 9.81$   $\mu\text{g/g dw}$ ), and Zn ( $253.34 \pm 22.76$   $\mu\text{g/g dw}$ ) of car park dust; Zn ( $997.20 \pm 16.10$   $\mu\text{g/g dw}$ ) (of ceiling fan dust)] and K10 [Ni ( $26.88 \pm 1.84$   $\mu\text{g/g dw}$ ) and Zn ( $199.77 \pm 6.64$   $\mu\text{g/g dw}$ ) of car park dust; Cu ( $468.55 \pm 3.67$   $\mu\text{g/g dw}$ ), Ni ( $83.96 \pm 0.75$   $\mu\text{g/g dw}$ ), and Fe ( $3131.58 \pm 27.01$   $\mu\text{g/g dw}$ ) of ceiling fan dust] exhibited elevated concentrations of heavy metals that might be related to vehicular activities as compared to K2 and KOSAS. In general, ceiling fan dust had significantly ( $P < 0.05$ ) higher concentrations of heavy metals than the car park dust. In comparison to other reported studies in the literature, the maximum levels of Ni and Cu were comparable or higher than those reported for major cities in the world. Hence, more monitoring studies should be conducted in the future to check for metal contamination in the dust, as this can serve as an atmospheric indicator of heavy metal pollution.

**Keywords:** Dust, heavy metals, UPM residential colleges

### INTRODUCTION

Dust is also known as a particulate matter that can range from 1 to 1000 $\mu\text{m}$  in size and it receives inputs from various urban sources (Meza-Figueroa *et al.*, 2007). Heavy metals in dust particulate form will eventually be deposited on the ground and can easily be suspended by wind into the atmosphere (Sharma *et al.*, 2008; Amato *et al.*, 2009). They are potentially being hazardous to human health, especially to children (Lin *et al.*, 2002; Ng *et al.*, 2003; Meza-Figueroa *et al.*, 2007). Ferreira-Baptista and Miguel (2005) define street dust as solid particles that comprise of impervious materials and accumulate outdoors in urban environments. Street or urban roadside dust consist of vehicular exhaust particles, household dust, soil dust, construction dust and aerosols that are carried freely by air and water (Takada *et al.*, 1991; Al-Khashman, 2004; Meza-Figueroa *et al.*, 2007). Atmospheric pollution is one of the major sources of heavy metal contamination

*al.*, 2003; Meza-Figueroa *et al.*, 2007). Ferreira-Baptista and Miguel (2005) define street dust as solid particles that comprise of impervious materials and accumulate outdoors in urban environments. Street or urban roadside dust consist of vehicular exhaust particles, household dust, soil dust, construction dust and aerosols that are carried freely by air and water (Takada *et al.*, 1991; Al-Khashman, 2004; Meza-Figueroa *et al.*, 2007). Atmospheric pollution is one of the major sources of heavy metal contamination

Received: 31 July 2009

Accepted: 25 January 2010

\*Corresponding Author

in soils and street dust in urban areas (Lin *et al.*, 2001). Although studies carried out on ceiling fan dust are rather few, there are more closely related research done on household or indoor dust that have mainly focused on floor or carpet dust (e.g. Kim & Fergusson, 1993; Tong & Lam, 1998; Jabeen *et al.*, 2001). Loading of heavy metals in the indoor environment are actually related to the outdoor environment (Kim & Fergusson, 1993; Tong & Lam, 1998; Jabeen *et al.*, 2001). Despite the abundant literature about dust, only Yap *et al.* (2007) have reported heavy metal levels in indoor ceiling fan dust in Ipoh and Serdang, Malaysia.

Since dust always presents in the urban environment and has the potential to contribute to metal toxicity in humans, there is a growing concern over this threat in the scientific community (Ng *et al.*, 2003). The US EPA (1986) reported that populations exposed to Cd, Cu, and Zn pollutants are in risk of developing nervous system alterations which can lead to health hazards. The emissions from roadways can cause respiratory illnesses (Lin *et al.*, 2002). There have been studies done on street dust near places where children play (e.g. Ng *et al.*, 2003; Meza-Figueroa *et al.*, 2007) since children are more sensitive to contaminant-bearing dust (Meza-Figueroa *et al.*, 2007).

The environmental impacts of atmospheric deposition have been studied for more than a century, and the first effect that was described on a scientific basis was probably the decline of epiphytic lichens in areas with high levels of atmospheric pollution (Wolterbeek, 2002). Heavy metals such as Cu, Pb, and Zn can be accumulated in soils from atmospheric deposition from sources such as vehicular exhausts, industrial discharges, oil lubricants, automobile parts, and corrosion of building materials (Li *et al.*, 2001). The determination of metal levels in environmental samples in top soils and urban roadside dust is therefore necessary for monitoring environmental pollution since such levels can affect the surrounding ecosystems and heavy metals can accumulate in organisms (Aksoy & Demirezen, 2006). Due to the significant contribution of street dust to pollution

in the urban environment, numerous studies on heavy metals of street dust have been carried out in developed countries (Charlesworth *et al.*, 2003; Manno *et al.*, 2006), but such studies are very much lacking in developing countries.

The objectives of this study were to determine the concentrations of metals on ceiling fan and car park dust collected from four different residential colleges in the Serdang campus of Universiti Putra Malaysia (UPM) and to relate their anthropogenic sources to the human activities observed in the surroundings.

## MATERIALS AND METHODS

Samples of dust were collected from four residential colleges in UPM, namely K2, K5, KOSAS, K10, in February 2009 (*Fig. 1*). The dust was collected from ceiling fans at each college, and sandy dust at car parks. For the ceiling fan dust, only the one in the rooms located on the first floor was collected. All the samples were collected from non-rusty ceiling fans. For the roadside car park dust, sand dust on top of the car park was collected using a polyethylene brush, and tray. The dust was stored in clean and labelled polyethylene plastic bags. For the preliminary study, three (3) sites at KOSAS, K2, and K5 were considered as the reference sites to K10 which is closest to main north-south highway. However, it was not certain if the metals levels would be the highest at K10, and it was for this reason that the present study was conducted.

After drying at 80°C in an oven until constant dry weights, the car park sand dust was sieved through a 53µm sieving tool. However, the fan dust samples were not sieved as they were processed immediately after drying. This is because all the airborne particles that can be trapped by fans are potentially inhaled by humans (Yap *et al.*, 2007).

The methodology of the analysis of heavy metals in the dust is similar to those described by Yap *et al.* (2002). In brief, the direct aqua-regia method was used to process the digested sample. About 0.5g of the dust sample was weighed and placed in digestion tube (3 replicates) for ceiling



TABLE 1  
Comparison of the mean concentrations ( $\mu\text{g/g}$  dry weight) of Cd, Cu, Fe, and Zn  
between ceiling fan dust and car park sand dust using the Mann-Whitney test

| Metal | College   | N  | Ceiling Fan Dust                  | N  | Car Park Dust                     | Mann-Whitney (p-value) |
|-------|-----------|----|-----------------------------------|----|-----------------------------------|------------------------|
| Cd    | K2        | 9  | 4.15 $\pm$ 0.84 <sup>A</sup>      | 3  | 3.01 $\pm$ 0.29 <sup>c</sup>      | P=0.518 (P>0.05)       |
|       | K5        | 6  | 3.84 $\pm$ 0.03 <sup>A</sup>      | 3  | 2.06 $\pm$ 0.04 <sup>ab</sup>     | P=0.020 (P<0.05)       |
|       | KOSAS     | 9  | 3.50 $\pm$ 0.25 <sup>A</sup>      | 3  | 2.19 $\pm$ 0.05 <sup>b</sup>      | P=0.013 (P<0.05)       |
|       | K10       | 15 | 3.58 $\pm$ 0.22 <sup>A</sup>      | 3  | 1.24 $\pm$ 0.08 <sup>a</sup>      | P=0.008 (P<0.05)       |
|       | Average   | 39 | 3.73 $\pm$ 0.17                   | 12 | 2.11 $\pm$ 0.12                   | P< 0.05                |
|       | (Min–Max) |    | (3.15-5.83)                       |    | (0.72-3.76)                       |                        |
| Zn    | K2        | 9  | 800.15 $\pm$ 8.44 <sup>A</sup>    | 3  | 103.68 $\pm$ 16.34 <sup>a</sup>   | P=0.013 (P<0.05)       |
|       | K5        | 6  | 997.20 $\pm$ 16.10 <sup>C</sup>   | 3  | 253.34 $\pm$ 22.76 <sup>c</sup>   | P=0.020 (P<0.05)       |
|       | KOSAS     | 9  | 850.24 $\pm$ 12.20 <sup>B</sup>   | 3  | 164.13 $\pm$ 5.65 <sup>b</sup>    | P=0.013 (P<0.05)       |
|       | K10       | 15 | 897.13 $\pm$ 20.74 <sup>B</sup>   | 3  | 199.77 $\pm$ 6.64 <sup>b</sup>    | P=0.008 (P<0.05)       |
|       | Average   | 39 | 892.11 $\pm$ 18.72                | 12 | 177.61 $\pm$ 9.68                 | (P< 0.05)              |
|       | (Min–Max) |    | (785.60-1029.08)                  |    | (52.27-304.64)                    |                        |
| Fe    | K2        | 9  | 2631.62 $\pm$ 14.83 <sup>A</sup>  | 3  | 1687.22 $\pm$ 37.88 <sup>a</sup>  | P=0.013 (P<0.05)       |
|       | K5        | 6  | 3008.97 $\pm$ 68.11 <sup>BC</sup> | 3  | 1802.40 $\pm$ 9.81 <sup>c</sup>   | P=0.020 (P<0.05)       |
|       | KOSAS     | 9  | 2894.41 $\pm$ 36.34 <sup>B</sup>  | 3  | 1691.61 $\pm$ 28.66 <sup>ab</sup> | P=0.013 (P<0.05)       |
|       | K10       | 15 | 3131.58 $\pm$ 27.01 <sup>C</sup>  | 3  | 1783.25 $\pm$ 5.40 <sup>bc</sup>  | P=0.008 (P<0.05)       |
|       | Average   | 39 | 2976.41 $\pm$ 56.36               | 12 | 1742.89 $\pm$ 13.47               | (P< 0.05)              |
|       | (Min–Max) |    | (2603.72-3243.56)                 |    | (1552.55-1846.71)                 |                        |
| Ni    | K2        | 9  | 46.64 $\pm$ 2.03 <sup>A</sup>     | 3  | 15.68 $\pm$ 1.21 <sup>a</sup>     | P=0.013 (P<0.05)       |
|       | K5        | 6  | 46.74 $\pm$ 0.46 <sup>A</sup>     | 3  | 19.66 $\pm$ 0.57 <sup>a</sup>     | P=0.020 (P<0.05)       |
|       | KOSAS     | 9  | 58.12 $\pm$ 2.23 <sup>B</sup>     | 3  | 13.80 $\pm$ 0.95 <sup>a</sup>     | P=0.013 (P<0.05)       |
|       | K10       | 15 | 83.96 $\pm$ 0.75 <sup>C</sup>     | 3  | 26.88 $\pm$ 1.84 <sup>b</sup>     | P=0.008 (P<0.05)       |
|       | Average   | 39 | 66.13 $\pm$ 5.35                  | 12 | 20.17 $\pm$ 1.19                  | (P< 0.05)              |
|       | (Min–Max) |    | (42.59-96.35)                     |    | (9.85-38.41)                      |                        |
| Cu    | K2        | 9  | 226.27 $\pm$ 1.23 <sup>A</sup>    | 3  | 18.95 $\pm$ 0.70 <sup>a</sup>     | P=0.013 (P<0.05)       |
|       | K5        | 6  | 277.29 $\pm$ 7.69 <sup>B</sup>    | 3  | 62.94 $\pm$ 0.77 <sup>c</sup>     | P=0.013 (P<0.05)       |
|       | KOSAS     | 9  | 304.22 $\pm$ 11.64 <sup>C</sup>   | 3  | 30.93 $\pm$ 4.08 <sup>ab</sup>    | P=0.013 (P<0.05)       |
|       | K10       | 15 | 468.55 $\pm$ 3.67 <sup>D</sup>    | 3  | 43.59 $\pm$ 3.99 <sup>b</sup>     | P=0.008 (P<0.05)       |
|       | Average   | 39 | 360.51 $\pm$ 31.11                | 12 | 37.96 $\pm$ 2.91                  | (P< 0.05)              |
|       | (Min–Max) |    | (223.84-536.00)                   |    | (16.11-76.46)                     |                        |

Note: N= numbers of samples analysed

Metal concentrations at the different sites sharing common letters (<sup>a</sup>, <sup>b</sup>, and <sup>c</sup>) are not significantly different for Gabriel, One-way ANOVA testing. (P<0.05)

Metal concentrations at the different sites sharing common capital letters (<sup>A</sup>, <sup>B</sup>, and <sup>C</sup>) are not significantly different for Student-Newman-Keuls, One-way ANOVA testing (P<0.05)

colleges, whereas K10 had the highest metal concentration of Ni ( $26.88 \pm 1.84$ ), and K2 had the highest concentration of Cd ( $3.01 \pm 0.29$ ). On the other hand, K2 was found to have the lowest concentrations of Zn ( $103.68 \pm 16.34$ ), Fe ( $1687.22 \pm 37.88$ ), while Cu ( $18.95 \pm 0.70$ ). KOSAS was found to have the lowest concentration of Ni ( $13.80 \pm 0.95$ ) and K10 had the lowest concentration of Cd ( $1.24 \pm 0.08$ ).

As for the ceiling fan dust, K2 was found to have the highest concentration of Cd ( $4.15 \pm 0.84$ ) as compared to other residential colleges. Meanwhile, K5 had the highest concentration of Zn ( $997.20 \pm 16.10$ ). The highest concentrations of Fe ( $83.96 \pm 0.75$ ), Ni ( $83.96 \pm 0.75$ ), and Cu ( $468.55 \pm 3.67$ ) were found in K10. Among all the residential colleges, the lowest concentrations of Cu ( $226.27 \pm 1.23$ ), Ni ( $46.64 \pm 2.03$ ), Fe ( $2631.62 \pm 14.83$ ), and Zn ( $800.15 \pm 8.44$ ) were found in K2, while KOSAS had the lowest concentration of Cd ( $3.50 \pm 0.25$ ).

Looking at the location of K5, the residential college's car park is situated next to a hill slope. Thus, higher amounts of fuel were used for motor vehicles (buses, cars, motor, etc.) in low gear to go up the hill in addition to frequent braking while going downhill. Furthermore, buses also stop frequently at the Putra Food Court which is located near to K5. Hence, higher heavy metal concentrations could be expected at this site, and this corresponded well with the highest concentrations of Cu, Fe and Zn found in the car park sand dust at K5 and the high Zn concentration in the ceiling fan dust of K5. Cu and Zn could be derived from the mechanical abrasion of vehicles as they are used in the production of brass alloy and they could also come from brake linings, oil leak sumps and cylinder head gaskets (Jiries *et al.*, 2001). Tyre to road friction would be higher for up slope and down slope driving and increased braking manoeuvres which contribute to the higher mechanical abrasion of vehicles and greater stop-start manoeuvres (Ellis & Revitt, 1982). However, K5 had significantly lower Ni concentration than K10 ( $p < 0.05$ , One-way ANOVA test), the residential college which is located nearest to the highways (Lebuhraya Utara-Selatan

and Lebuhraya Sungai Besi). There is no doubt that car fuel combustion is higher (with higher volume of cars on the highway) along the vicinity of the highways. Thus, this could explain the high concentration of Ni in the car park dust and the high concentrations of Cu, Ni, and Zn in the ceiling fan dust. In particular, Cu is a common element that is used in automobile thrust bearing, brake lining, and other parts of the engine (Ng *et al.*, 2003). Zn may also originate from lubricant oil and tyres of motor vehicles (Meza-Figueroa *et al.*, 2007). Ni could be involved in the vehicular fuel combustion process (Meza-Figueroa *et al.*, 2007). However, the ceiling fan dust collected in K10 had a significantly ( $p < 0.05$ ) higher concentration of Fe as well. Fe is often the major component of brake pads (Amato *et al.*, 2009). This could be due to the outdoor sources of Fe contributing to the room dust, as the residential college is situated next to the highway. The concentrations of heavy metal were reported to be higher on streets where traffic was more likely to undergo stop-start manoeuvres (Ellis & Revitt, 1982).

In general, the accumulation of metal found in the ceiling fan dust was observed to be resulted from different factors such as indoor pollutant sources, infiltration of outdoor dust, and the absorption of metals due to indoor humidity (Davies *et al.*, 1987). Most of the rooms where the samples were taken lack proper ventilation as the windows were either seldom or never opened by the occupants. This is in line with study by Jabeen *et al.* (2001) who showed that the interior house dust had higher concentrations of Cd and Pb, even for houses with little ventilation. Furthermore, the ceiling fan dust collected in this study had been there for a long time as the fans were not cleaned for several months to years. Therefore, heavy metal concentrations may have increased with time as compared to the outdoor street dust, as the conditions of the interior are much more stable than the outdoors.

In addition, weather is likely to influence the heavy metal distribution, concentration and retention time of the deposited street dust (Charlesworth *et al.*, 2003). A study by

Fergusson *et al.* (1984) showed that metal loadings for Cd, Cu, Pb, and Zn in house dust correlated strongly with the amount of dust in the house and this in turn correlated strongly to the amount of carpet used (Jabeen *et al.*, 2001).

K2 was found to have significantly ( $p < 0.05$ ) lower levels of Cu, Fe, Ni, and Zn for its car park dust, while Cu, Ni, Fe, and Zn ceiling fan dust when compared to the other residential colleges. KOSAS had significantly ( $p < 0.05$ ) lower concentrations of Cu, Ni, and Zn for the car park dust, and Cu and Zn for ceiling fan dust relative to the other residential colleges. The lower metal concentrations found in these two colleges could be due to the lesser road traffic activities around there as compared to K10 and K5. However, K2 had a significantly ( $p < 0.05$ ) higher concentration of Cd for its car park dust, and this was also higher but insignificant for the ceiling fan dust as compared to K10.

The correlations between heavy metal concentrations among all the car park sand dust are presented in Table 2. Based on the data given, significant (at least  $p < 0.05$ ) and positive correlations were found between Zn-Fe (0.320), Zn-Ni (0.502), Zn-Cu (0.625), Ni-Fe (0.323), and Cu-Fe (0.580). However, the positive relationships were found to be not convincing, and thus, further investigations are needed. As for the ceiling fan dust (Table 3), significant (at least  $p < 0.05$ ) and positive correlations were found between Fe-Ni (0.732), Fe-Cu (0.846), and Cu-Ni (0.918). The strong and positive correlations among Cu, Fe, Ni, and Zn in the car park dust, as well as Cu, Fe, and Ni in the ceiling fan dust might indicate that they could have come from the same sources, such as road traffic, as these metals are contained in motor vehicles. The study by Al-Khashman (2004) revealed a high correlation of the metal concentrations of Cu, Fe, Pb, and Zn which convinced him that anthropogenic activities are the main sources of heavy metal in soils. Table 3 shows that a larger proportion of Cd might have come from different sources, even though it could also be due to road traffic activities, as reported by Meza-Figueroa *et al.* (2007). This could explain why K2 had a significant ( $p < 0.05$ ) higher concentration of

Cd than K10. Cd might have come from colour pigments, like the ones from carpets (Jabeen *et al.*, 2001). Cd could also come from air-borne emissions from various sources (Barnejee, 2003), as it is used to protect the surface of brass from corrosion (Charlesworth *et al.*, 2003).

TABLE 2  
Pearson's Correlations for the car park dust (N=39)

|    | Cd | Zn      | Fe       | Ni       | Cu      |
|----|----|---------|----------|----------|---------|
| Cd | 1  | -0.404* | -0.747** | -0.439** | -0.371* |
| Zn |    | 1       | 0.320*   | 0.502**  | 0.625** |
| Fe |    |         | 1        | 0.323*   | 0.580** |
| Ni |    |         |          | 1        | 0.279   |
| Cu |    |         |          |          | 1       |

TABLE 3  
Spearman's correlations for the ceiling fan dust (N=15)

|    | Cd | Zn    | Fe      | Ni      | Cu      |
|----|----|-------|---------|---------|---------|
| Cd | 1  | 0.064 | -0.018  | -0.129  | 0.007   |
| Zn |    | 1     | 0.664** | 0.089   | 0.407   |
| Fe |    |       | 1       | 0.732** | 0.846** |
| Ni |    |       |         | 1       | 0.918** |
| Cu |    |       |         |         | 1       |

Note: \*= Correlation is significant at the 0.05 level (2-tailed).  
\*\*= Correlation is significant at the 0.01 level (2-tailed).

When compared to other related studies in the literature, the mean Zn concentration for the car park dust in the present study (Table 4) was found to be higher than that of the street dust in Bahrain (Akhter & Madany, 1993). However, the concentrations of Cd, Cu, and Ni found in the car park dust in this study were lower as compared to the street dust from other studies (e.g. as reported by Fergusson & Ryan, 1984; Lehame *et al.*, 1992; Akhter & Madany, 1993; Chon *et al.*, 1995; Wang *et al.*, 1998).

Table 5 shows that the overall maximum concentration ranges of Cu, Ni, and Zn in the ceiling fan dust were higher than those reported by Yap *et al.* (2007), and these increase the concern over their influence on human health. On the contrary, the overall concentrations of Cd and Fe in the ceiling fan dust were lower than those reported by Yap *et al.* (2007). Fe

TABLE 4  
Comparison of the results from the global studies on heavy metal concentrations in street dust ( $\mu\text{g/g}$ ), as compiled by Charlesworth *et al.* (2003) and in this study

| City      | Reference                   | Metal     |             |            |              |
|-----------|-----------------------------|-----------|-------------|------------|--------------|
|           |                             | Cd        | Cu          | Ni         | Zn           |
| New York  | Fergusson & Ryan (1984)     | 8         | 355         | -          | 1811         |
| Seoul     | Chon <i>et al.</i> (1995)   | 3         | 101         | -          | 296          |
| London    | Lehame <i>et al.</i> (1992) | 6250      | 61-323      | 32-74      | -            |
| Hong Kong | Wang <i>et al.</i> (1998)   | -         | 92-392      | -          | 574-2397     |
| Bahrain   | Akhter & Madany (1993)      | 72        | -           | 126        | 152          |
| Serdang   | This study                  | 0.72-3.76 | 16.11-76.46 | 9.85-38.41 | 52.27-304.64 |

- indicates data are available

TABLE 5  
Comparisons for the values (minimum-maximum) of heavy metal concentrations ( $\mu\text{g/g}$  dry weight) in the fan dust collected in this study and those reported by Yap *et al.* (2007)

| Location             | Cd,<br>$\mu\text{g/g}$ | Zn,<br>$\mu\text{g/g}$ | Fe,<br>$\mu\text{g/g}$ | Ni,<br>$\mu\text{g/g}$ | Cu,<br>$\mu\text{g/g}$ | Reference                   |
|----------------------|------------------------|------------------------|------------------------|------------------------|------------------------|-----------------------------|
| Serdang,<br>Malaysia | 3.15-<br>5.83          | 785.60-<br>1029.08     | 2603.72-<br>3243.56    | 42.59-<br>96.35        | 223.84-<br>536.00      | This study                  |
| Ipoh, Malaysia       | 5.37-<br>16.70         | 563.20-<br>815.90      | 3470.70-<br>4455.46    | 43.97-<br>58.55        | 159.99-<br>229.32      | Yap <i>et al.</i><br>(2007) |
| Serdang,<br>Malaysia | 11.40-<br>13.71        | 688.39-<br>868.10      | 3446.50-<br>4440.39    | 28.51-<br>36.22        | 163.54-<br>270.05      | Yap <i>et al.</i><br>(2007) |

is a common element of the earth's crust, so its levels in the car park dust could be related to the mobilization of soil particles (Fung & Wong, 1995).

### CONCLUSION

In conclusion, K5 (Zn, Fe, and Cu, car park dust; Zn, ceiling fan dust) and K10 (Zn and Ni, car park dust; Cu, Ni, and Fe, ceiling fan dust) were found to have high concentrations of heavy metals, since they are comparable to other reported data for major cities, and this could be related to the vehicular traffic at these residential colleges. In general, ceiling fan dust had significantly ( $P < 0.05$ ) higher concentrations of heavy metals compared to that in the car park dust. Meanwhile, most of the heavy metals concentrations (Cd, Cu, Ni, and Fe) found in the

car park dust were still low than the data reported in the literature. However, the concentrations of Zn in the car park dust and the concentrations of Cu, Ni and Zn in the ceiling fan dust undertaken in this study were found to be comparable to or higher than those reported for other cities in the literature. Thus, further monitoring studies are needed since dust is considered as a potential atmospheric indicator of heavy metal pollution and its relationship to human health should be considered seriously.

### ACKNOWLEDGEMENTS

This work was financially supported by a Fundamental Research Grant (FRGS) 2007 provided by the Ministry of Higher Education, Malaysia, through Vote no. 5523266.

## REFERENCES

- Akhter, M. S., & Madany, I. M. (1993). Heavy metals in street and house dust in Bahrain. *Water, Air and Soil Pollution*, 66, 111 – 119.
- Aksoy, A., & Demirezen, D. (2006). *Fraxinus excelsior* as a biomonitor of heavy metal pollution. *Polish Journal of Environmental Studies*, 15, 27-33.
- Al-Khashman, O. A., (2004). Heavy metal distribution in dust, street dust and soils from the work place in Karak Industrial Estate, Jordan. *Atmospheric Environment*, 38, 6803-6812.
- Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A., & Moreno, T. (2009). Spatial and chemical patterns of PM10 in road dust deposited in urban environment. *Atmospheric Environment*, 43, 1650-1659.
- Banerjee, A. (2003). Heavy metal levels and solid phase speciation in street dust of Delhi, India. *Environmental Pollution*, 123, 95-105.
- Charlesworth, S., Everett, M., Mc Carthy, R., Ordonez, A., & De Minguel, E. (2003). A comparative study of heavy metal concentration and distribution in deposited street dust in large and a small urban area: Birmingham and Coventry, West Midlands, UK. *Environmental International*, 29, 563-573.
- Chon, H-T., Kim, K-W., & Kim, J-Y. (1995). Metal contamination of soils and dust in Seoul metropolitan city, Korea. *Environmental Geochemistry and Health*, 17, 139–146.
- Davies, D. J. A., Watt, J. M., & Thornton, I. (1987). Lead levels in Birmingham dust and soils. *Science of the Total Environment*, 67, 177– 85.
- Ellis, J. B., & Revitt, D. M., (1982). Incidence of heavy metals in street surface sediments: solubility and grain size studies. *Water, Air, and Soil Pollution*, 17, 87-100.
- Fergusson, J. E., & Ryan, D. E. (1984). The elemental composition of street dust from large and small urban areas related to city type, source and particle size. *Science of the Total Environment*, 34, 101– 116.
- Ferreira-Baptista, L., & De Miguel, E. (2005). Geochemistry and risk assessment of street dust in Luanda, Angola: A tropical urban environment. *Atmospheric Environment*, 39, 4501-4512.
- Fung, Y. S., & Wong, L. W. Y. (1995). Apportionment of air pollution sources by receptor models in Hong Kong. *Atmosphere Environment*, 29, 2041-2048.
- Jabeen, N. A., Ahmed, S., Hassan, S. T., & Alam, N. M. (2001). Levels and sources of heavy metals in house dust. *Journal of Radioanalytical and Nuclear Chemistry*, 247, 145-149.
- Jiries, A. G., Hussein, H. H., & Halas, Z., (2001). The quality of water and sediments of street runoff in Amman, Jordan. *Hydrology Proceedings*, 15, 815-824.
- Kim, N., & Fergusson, J. (1993). Concentrations and sources of cadmium, copper, lead, and zinc in house dust in Christchurch, New Zealand. *Science of the Total Environment*, 138, 1-21.
- Leharne, S., Charlesworth, D., & Choudhry, B. (1992). A survey of metal levels in street dust in an inner London neighbourhood. *Environment International*, 18, 263–270.
- Li, X., Poon, C. S., & Liu, P. S., (2001). Heavy metal contamination of urban soils and street dust in Hong Kong. *Applied Geochemistry*, 16, 1361-1368.
- Lin, S., Munsie, J. P., Hwang, S.-A., Fitzgerald, E., & Cayo, M. R. (2002). Childhood asthma hospitalization and residential exposure to state route traffic. *Environmental Research*, 88, 73-81.
- Manno, E., Varrica, D., & Dongarra, G. (2006). Metal distribution in road dust samples collected in an urban area close to a petrochemical plant at Gela, Sicily, Italy. *Atmospheric Environment*, 40, 5929 – 5941.
- Meza-Figueroa, D., De La O-Villanueva, M., & De La Parra, M. L. (2007). Heavy metal distribution in dust from elementary schools in Hermosillo, Sonora, Mexico. *Atmospheric Environment*, 41, 276-288.
- Ng, S. L., Chan, L. S., Lam, K. C., & Chan, W. K. (2003). Heavy metal contents and magnetic properties of playground dust in Hong Kong. *Environmental Monitoring and Assessment*, 89, 221–232.



- Sezgin, N., Ozcan, H. K., Demir, G., Nemlioglu, S., & Bayat, Cuma. (2003). Determination of heavy metal concentrations in street dust in Istanbul E-5 highway. *Environment International*, 29, 979-985.
- Sharma, R. K., Agrawal, M., & Marshall, F. M. (2008). Atmospheric deposition of heavy metals (Cu, Zn, Cd and Pb) in Varanasi City, India. *Environment Monitoring and Assessment*, 142, 269–278.
- Takada, H., Honda, T., Harada, M., & Ogura, N. (1991). Distribution and sources of polycyclic aromatic hydrocarbons (PAHs) in street dust from the Tokyo Metropolitan area. *Science of Total Environment*, 107, 45-69.
- Tong S. T. Y. (1998). Indoor and outdoor household dust contamination in Cincinnati, Ohio, USA. *Environmental Geochemistry and Health*, 20, 123– 133.
- Tong, S. T. Y., & Lam, K. C. (1998). Are nursery schools and kindergartens safe for our kids? The Hong Kong study. *Science of Total Environment*, 216, 217-225.
- U.S. EPA (US Environment Protection Agency), (1986). *Air Quality Criteria for Mercury and Lead*. EPA. Research Triangle Park, NC.
- Wang, W. H., Wong, M. H., Leharne, S., & Fisher, B. (1998). Fractionation and biotoxicity of heavy metals in urban dust collected from Hong Kong and London. *Environmental Geochemistry and Health*, 20, 195– 198.
- Wolterbeek, B. (2002). Biomonitoring of trace element air pollution: principles, possibilities and perspectives. *Environmental Pollution*, 120, 11-21.
- Yap, C. K., Ismail, A., & Tan, S. G. (2007). Heavy metal concentrations in indoor fan dust of residential areas: A preliminary study. *Malaysian Applied Biology*, 36, 47-49.
- Yap, C. K., Ismail, A., Tan, S. G., & Omar, H., (2002). Correlations between speciation of Cd, Cu, Pb and Zn in sediment and their concentrations in total soft tissue of green-lipped mussel *Perna viridis* from the west coast of Peninsular Malaysia. *Environment International*, 28, 117-126.